WK	S
Mass-Mass Ca	alculations

NAME		
Period _	Date	

Perform the following stoichiometry calculations. Show your work, make sure all numbers have units, and watch Sig Figs! Remember:

$$Mass \rightarrow Moles \rightarrow Moles \rightarrow Mass$$

1) Titanium is a transition metal used in many alloys because it is extremely strong and lightweight. Titanium(IV) chloride (TiCl₄) is extracted from titanium oxide using chlorine and coke (carbon):

$$TiO_2(s) + C(s) + 2 Cl_2(g) \rightarrow TiCl_4(s) + CO_2(g)$$

If you begin with 1.25 mol TiO₂, what mass of Cl₂ gas is needed?

2) At high temperature, molten sodium chloride is decomposed into the elements sodium (a liquid at these temperatures) and chlorine (a gas) by means of electrical energy. How many grams of chlorine gas can be obtained from 2.50 mol NaCl? (You need to write and balance the chemical equation!)

3) Aluminum oxidizes according the following unbalanced equation:

$$Al(s) + O_2(g) \rightarrow Al_2O_3(s)$$

- a. Balance the equation
- b. How many grams of Al₂O₃ would be formed by the reaction of 29.75 grams of Al with enough O₂?

c. How many grams of O₂ would be required to form 65.32 grams of Al₂O₃?

4)		nitrogen pentoxide is an acidic gas that reacts with water to forms aqueous nitric acid. Write and balance the chemical equation.
	b.	How many grams of dinitrogen pentoxide would be required to completely react with 13.44 grams of water?
	c.	How many grams of dinitrogen pentoxide would be needed, given enough water, to produce 105.65 grams of nitric acid?
5)	car	lid iron(III) oxide reacts with gaseous carbon monoxide at high temperature and forms iron metal and bon dioxide gas. Write and balance the chemical equation.
	b.	How many grams of carbon dioxide would be formed in the reaction where 25.1 grams of iron are formed?
	c.	How many grams of iron would be produced by the reaction of 182.0 grams of iron(III) oxide with sufficient carbon monoxide?
	; ⁷ O	Answers: 1) 177 g Cl ₂ ; 2) 88.6 g Cl ₂ ; 3b) 56.18 g Al ₂ O ₃ ; 3c) 30.75 g O ₂ ; 4b) 80.56 g N ₂ O ₅ ; 4c) 90.55 g N ₂ O ₅ ; 5b) 29.7 g C (5c) 127.3 g Fe