

3.104 The carat is a unit of mass used by jewelers. One carat is exactly 200 mg. How many carbon atoms are in a 24-carat diamond?

3.107 An impure sample of Zn is treated with an excess of H_2SO_4 to form ZnSO_4 and H_2 .
(a) Write the balanced equation for the reaction.

(b) If 0.0764 g of H_2 is obtained from 3.86 g of Zn, calculate the % purity of the Zn.

(c) What assumptions must you make in (b)?

3.131 The formula of a hydrate of barium chloride is $\text{BaCl}_2 \cdot x\text{H}_2\text{O}$. If 1.936 g of the hydrate gives 1.864 g of BaSO_4 upon treatment with sulfuric acid, calculate the value of x .

(HINT: $\text{BaCl}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + 2 \text{HCl}$)

3.134 When 0.273 g of Mg is heated strongly in an N_2 atmosphere, a chemical reaction occurs. The product has a mass of 0.378 g. Calculate the empirical formula of the compound, containing only Mg and N and name it.

3.136 The anti-knock additive in leaded gasoline contains only C, H, and Pb. When 51.36 g of this compound is burned in an apparatus like that in Figure 3.6, 55.90 g of CO₂ and 28.61 g of H₂O are produced. Determine the empirical formula of the additive.

Additional Questions:

A. 0.755 g sample of hydrated copper(II) sulfate is heated carefully until it had changed completely to anhydrous copper(II) sulfate with a mass of 0.483 g. Determine the value of x in the formula of the hydrate, CuSO₄ · x H₂O. What would you have done in the lab to be sure that no water was left in the sample after heating?

B. Write the balanced chemical equation for the following reactions:


- i. calcium metal reacts with water to produce aqueous calcium hydroxide and hydrogen gas.

- ii. barium hydroxide reacts with sulfuric acid to produce barium sulfate and water.

- iii. iron(III) sulfide reacts with hydrogen chloride to form iron(III) chloride and hydrogen sulfide.

- iv. carbon disulfide reacts with ammonia to produce hydrogen sulfide and ammonium thiocyanate.

C. Hydrogen peroxide can be produced by the following reaction:



i. What is the theoretical yield (in grams) of hydrogen peroxide when 1.50g of barium peroxide is treated with 25.0 mL of a hydrochloric acid solution containing 0.0272g of HCl per mL?

ii. How many moles of the excess reactant are left unreacted?

D. Methyl isothiocyanate (MITC), an organosulfur compound which contains only C, H, N, and S, is used in agriculture as a soil fumigant, mainly for protection against fungi and nematodes. Find the empirical formula for MITC if combustion analysis of a 0.2415-g sample gives 0.2907 g CO₂, 0.08926 g H₂O, a mixture of nitrogen oxides, and 0.2116 g SO₂.

Answers: 104) 2.4×10^{23} atoms C; 107b) 64.2%; 131) BaCl₂ · 2H₂O; 134) Mg₃N₂; 136) PbC₈H₂₀; A) $\frac{5}{2}$; C) 0.298 g H₂O₂; C) 0.010 mol; D) C₂H₃NS